

**REVIEW PAPER**

**BIOCHEMISTRY OF MAGNESIUM**

**Kazimierz Pasternak, Joanna Kocot, Anna Horecka**

**Chair and Department of Medical Chemistry  
Medical University of Lublin**

Abstract

Magnesium is essential for biochemical functions of cells. Since  $\text{Mg}^{2+}$  has a relatively low ionic radius in proportion to the size of the nucleus (0.86 versus 1.14 f Å for  $\text{Ca}^{2+}$ ), it shows exceptional biochemical activity. Due to its physicochemical properties, intracellular magnesium can bind to the nucleus, ribosomes, cell membranes or macromolecules occurring in the cell's cytosol. It is indispensable for the nucleus to function as a whole and for the maintenance of physical stability as well as aggregation of ribosomes into polysomes able to initiate protein synthesis.  $\text{Mg}^{2+}$  can also act as a cofactor for ribonucleic acid enzymes (ribozymes) capable of specifically recognizing and cleaving the target mRNA. As an essential cofactor in NER, BER, MMR processes,  $\text{Mg}^{2+}$  is required for the removal of DNA damage. An activator of over 300 different enzymes, magnesium participates in many metabolic processes, such as glycolysis, Krebs cycle,  $\beta$ -oxidation or ion transport across cell membranes.  $\text{Mg}^{2+}$  plays a key role in the regulation of functions of mitochondria, including the control of their volume, composition of ions and ATP production.

**Key words:** magnesium, DNA repair process, enzyme, metabolic cycle, cellular respiration, calcium ion transport, potassium ion transport.

## BIOCHEMIA MAGNEZU

### Abstrakt

Magnez jest składnikiem niezbędnym dla zasadniczych funkcji biochemicznych komórki. Ponieważ  $Mg^{2+}$  ma relatywnie mały promień w stosunku do wymiarów jądra (0.86 i 1.14 Å odpowiednio dla  $Mg^{2+}$  i  $Ca^{2+}$ ), wykazuje dużą aktywność biochemiczną. Dzięki właściwościom fizykochemicznym śródkomórkowy  $Mg^{2+}$  może wiązać się z jądrem komórkowym, rybosomami, błonami komórkowymi oraz makromolekułami cytosolu komórki. Magnez jest niezbędny dla funkcjonowania jądra komórkowego jako całości oraz utrzymania fizycznej stabilności i agregacji rybosomów do polisomów zdolnych do biosyntezy białka. Odgrywa on również rolę kofaktora katalitycznych cząsteczek RNA (rybozymów), odpowiedzialnych za specyficzne rozpoznawanie i fragmentację docelowego mRNA. Jako kofaktor w procesach: NER, BER, MMR, przyczynia się do usuwania uszkodzeń DNA. Magnez, będąc aktywatorem ponad 300 różnych enzymów, uczestniczy w przebiegu wielu szlaków metabolicznych, takich jak glikoliza, cykl Krebsa,  $\beta$ -oksydacja czy transport jonów poprzez błony komórkowe. Odgrywa on ponadto bardzo ważną rolę w regulowaniu funkcji mitochondriów, łącznie z regulacją ich wielkości, kompozycją jonów, a także bioenergetyką i regulacją produkcji ATP.

Słowa kluczowe: magnez, proces naprawy DNA, enzym, cykl metaboliczny, oddychanie wewnątrzkomórkowe, transport jonów wapnia, transport jonów potasu.

## INTRODUCTION

The involvement of magnesium ions ( $Mg^{2+}$ ) in metabolic processes is governed not only by their abundance in nature or relative amount in living organisms but also by their physicochemical characteristics.

Since  $Mg^{2+}$  has a relatively low ionic radius in proportion to the size of the nucleus (0.86 versus 1.14 f Å for Ca), it shows exceptional biochemical activity. Ionized  $Mg^{2+}$  usually coordinates with 6-7 molecules of  $H_2O$ , as in the case of  $MgCl_2 \cdot 6 H_2O$  or  $Mg_2SO_4 \cdot 7 H_2O$ , while Ca and Ba combine with 1 or 2 mols of  $H_2O$  ( $BaCl_2$  and  $CaCl_2$  respectively). In comparison to calcium (Ca), the most abundant cation in the human body,  $Mg^{2+}$  displays higher affinity for oxygen donor ligands, that is negatively charged carboxylates and phosphates or enolate moieties (WOLF, CITTADINI 2003). Mg-water coordination occurs in a typical octahedral conformation and thereby magnesium exhibits slower water exchange than other ions. Consequently, it is much bigger and more stable in comparison to Ca in biological systems (WOLF, CITTADINI 2003, WEDEKIND et al. 1995)

Considering stereochemical properties, nickel (Ni) most closely resembles  $Mg^{2+}$  (it has atoms of the same size of and identical water exchange constant). However,  $Ni^{2+}$  cannot compete with  $Mg^{2+}$  in living organisms both due to its paucity and tendency to bind nitrogen rather than oxygen.

At the cellular level, magnesium ions compete not only with Ca but also with protons or amines ( $-NH_2^+$ ). Protons are usually present in concen-





Fig. 1. Octahedral magnesium complexes (WOLF, CITTADINI 2003)

trations below  $10^{-7}$  M at pH 7 and link up to phosphate groups with a pKa of 6.5, which is significantly lower than that of Mg-phosphate complexes. This suggests that  $Mg^{2+}$  is removed from ATP when pH falls to 6.0, causing significant modifications of Mg-dependent reactions:



Polyamines, organic derivatives of ammonia, exhibit high-affinity binding of polyanions, for instance nucleic acids, and dislodge  $Mg^{2+}$  bound therein (WOLF, CITTADINI 2003).

Due to its physicochemical properties, intracellular magnesium can bind to the nucleus, ribosomes, cell membranes or macromolecules occurring in the cell's cytosol.

## Magnesium and DNA

More than half the magnesium contained in the nucleus is closely associated with nucleic acids and free nucleotides. Since nucleic acids are polyanions, they require counterions in order to neutralize negatively charged phosphate groups (WOLF, CITTADINI 2003, ANASTASSOPOULOU, THEOPHANIDES 2002). The intracellular concentrations of Na and Ca are low, therefore the binding of the metal with nucleic acids is dominated by  $K^+$  and  $Mg^{2+}$ . Free  $Mg^{2+}$  is the winner in this competition because it has more positive charges (+ II and +I for Mg and K, respectively) and higher hydration energy (WOLF, CITTADINI 2003).

In Mg-DNA interactions, metal ions interact with purine bases at the N7 site and pyrimidine bases at the N3 site by forming chemical bonds,

Mg-N7 and/or Mg-N3. They also interact with the negatively charged oxygen atoms of phosphate groups of nucleotide chains (ANASTASSOPOULOU, THEOPHANIDES 2002). These interactions play a significant role in the stabilization of the secondary and tertiary structure of DNA.

The pathway of binding divalent metal ions to guanosino-5-monophosphate is shown in Figure 2. The intrinsic structure of this complex results from the fact that one of the coordinated water molecules may be substituted by the N7 coordination site of the nucleotide or be hydrogen-bound to it, while another one may be involved in a hydrogen bond with O6, and yet others form more hydrogen bonds (ANASTASSOPOULOU, THEOPHANIDES 2002).





Fig. 2. Structure of magnesium hydrate complex with guanosino-5-monophosphate (the broken lines show hydrogen bonds together with the hydrogen bond distances and angles) (ANASTASSOPOULOU, THEOPHANIDES 2002)

Whether or not  $Mg^{2+}$  acts as a gene regulator remains unclear. Since the bioactivity of this cation is remarkable, it is reasonable to think that magnesium may act as a competitor to polyamines, which are currently recognized as potential regulators of the cell cycle (WOLF, CITTADINI 2003).



Magnesium ions can affect the cellular cycle also in the form of Mg-ATP. This complex plays a key part in a phosphorylation cascade catalyzed by protein kinases. Alternatively, due to its capability to directly interact with proteins,  $Mg^{2+}$  can modulate histone phosphorylation (WOLF, CITTADINI 2003).

Irrespective of the above mechanisms, magnesium is indispensable for the functioning of the cell nucleus as a whole as it is involved in the activation of enzymes important for DNA repair (endonuclease) (WOLF, CITTADINI 2003, HARTWIG 2001, WOLF et al. 2003), replication (topoisomerase II (WOLF,

CITTADINI 2003, WOLF et al. 2003), polimerase I (WOLF, CITTADINI 2003) and transcription (rybonuclease H) (WOLF, CITTADINI 2003).  $Mg^{2+}$  is crucial for the physical integrality of double-stranded DNA.

In ribosomes,  $Mg^{2+}$  is associated with rRNA or proteins, which are essential for the maintenance of physical stability as well as aggregation of these structures into polysomes able to initiate protein synthesis. Cowan has shown that magnesium deficit leads to the cleaving of a ribosomal complex (COWAN 1995). Since the only function performed by ribosomes is protein biosynthesis, it is the presence of  $Mg^{2+}$  in ribosomes that conditions the shape of RNA structures by stimulating the transformation of amino acids into active forms, polypeptide synthesis and stabilization of a protein structure.

Moreover, magnesium can also act as a cofactor for ribonucleic acid enzymes (ribozymes) capable of specifically recognizing and cleaving the target mRNA. Ribozymes are chiefly used in steered therapy of neoplastic diseases (JOŠKO, KNEFEL 2003). It is believed that two metal ions (mostly  $Mg^{2+}$ , although  $Mn^{2+}$ ,  $Zn^{2+}$ ,  $Ca^{2+}$ ,  $Co^{2+}$  or  $Na^+$  are also possible) are necessary for catalytic activity of hammerhead ribozymes. One metal ion activates the attacking hydroxyl group, and the other stabilizes the negative charge of the oxygen atom of the released group (ADAMALA, PIKUŁA 2004). While one experiment with minimal hammerhead domains has demonstrated that the efficiency of catalysis is highly dependent on the concentration of magnesium ions, another one has shown that this efficiency can be increased at low magnesium concentration through stabilization of catalytically active conformation by tertiary interactions between helices I and II. Apart from these electrostatic interactions, both free  $Mg^{2+}$  as well as the GTP-Mg complex play an important role in tubulin polymerization and, consequently, in chromosome segregation during mitosis (HARTWIG 2001).

### **Role of magnesium in genomic stability**

In 1976 LOEB et al. noted that magnesium ions are indispensable for DNA replication fidelity. Although  $Co^{2+}$ ,  $Mn^{2+}$  and  $Ni^{2+}$  ions can be substituted for  $Mg^{2+}$ , such an exchange causes a considerable decrease in the fidelity of the discussed process. A metal ion (A) binds to the 3'-hydroxyl group of a new synthesis strand, leading to the lowering of its pKa and thereby facilitating an attack on the  $\alpha$ -phosphate of "arriving" dNTP. Another metal ion (B) facilitates the leaving of the  $\alpha$ - and  $\beta$ -phosphates as well as phosphodiester bond formation (HARTWIG 2001).

### **Role of magnesium in DNA repair processes**

Damage to DNA can be caused by exogenous factors (e.g. ultraviolet or electromagnetic radiation, high temperature, viruses, polycyclic aromatic hydrocarbons, radiotherapy or chemotherapy) or endogenous factors (mainly

ROS – reactive oxygen species). In order to lower the frequency of mutation, cells have developed many different DNA-repair systems.

Nucleotide excision repair (NER) is an evolutionarily conserved DNA repair pathway, which repairs DNA damaged by various environmental mutagens. Photodimers, pyrimidine, adducts as well as some of the damage repaired in the course of base excision repair (BER) can be removed from DNA (SANCAR 1994). The repair process is dependent on coordinated action of more than 20 different proteins. The majority of them are engaged in damage recognition and incision at both sides of the defect. Magnesium acts as a cofactor practically at every NER stage. Results of *in vitro* investigations have shown that this mechanism is completely inhibited in the case of absence and very high concentrations of magnesium. Experimental data have confirmed that the DNA-damage recognition protein UV-DDBP, the helicase XPD and the nuclease XPG are all magnesium-dependent. The element is required not only in enzymatic incision of DNA, but also in the processes of polymerization and ligation (HARTWIG 2001).

Reactive oxygen forms, normal products of cellular metabolism, lead to a wide variety of DNA modifications: destabilization of a DNA helix as well as degradation of protein-DNA crosslinks. Additionally, they are a major contributor to the oxidation of purine and pyrimidine bases, the damage of pentose ring, the hydrolysis of amine – or N-glycosidic bonds and phosphodiester bonds, hydrolytic deamination and methylation of oxygen or nitrogen atoms of DNA bases. At the extracellular level, ROS impair the function of blood platelets and induce protein, lipid or nucleic acid oxidation resulting in tissue destruction in many organs (CERIELLO, MOTZ 2004). It is believed that a mature organism can produce about 2 kilograms of superoxide anions per year, which can be transformed to  $H_2O_2$  by dismutation reaction (ROSZKOWSKI 2002).

Endogenous damages of DNA are mainly repaired by the BER mechanism. According to current models, BER begins with a removal of modified nitrogenous base by a specific N-glycosidase generating an AP site, which is repaired by AP-endonucleases cleaving the phosphodiester bond at the AP 5' side and leaving a 3' hydroxy terminus, making the action of DNA polymerase and ligase possible (ROSZKOWSKI 2002). Contrary to DNA glycosidases, enzymes involved in later stages of BER always require magnesium. In hydrolytic nucleases, which are metal-ion-dependent (mainly magnesium-dependent), metal interacts with a substrate or is directly involved in cleavage of the phosphate-oxygen bond. In human apurinic/aprimidinic endonuclease (HAP1), single magnesium ion combines with a defined Glu residue in the active center and aids the attack on the P-O3' bond by polarization of the P-O bond, perhaps by correctly orientating the phosphate group rather than directly participating in the nucleolysis reaction. HAP can also be activated by manganese or nickel ions, but its activity is considerably lower (by 50 and 90% respectively). Other examples of magnesium-dependent endonucle-

ases in BER include apurinic/apyrimidinic endonuclease, whose activity is associated with the 5-hydroxymethyluracil-DNA glycosylase, flap-endonuclease-1 (FEN-1) and a structure-specific endonuclease involved in DNA replication and DNA repair (HARTWIG 2001).

The third system, mismatch repair (MMR), has evolved to correct errors occurring during DNA replication or recombination of genes. Impairment of MMR leads to genomic instability, which creates favourable conditions for induction and development of carcinogenic processes. The most frequent mutations in the MMR system are those of genes from the MLH gene family. Constitutive mutations involving one allele of the MLH1 gene lead to tumor predisposition syndrome – hereditary nonpolyposis colorectal cancer (HNPCC type II) (COOK 2000) and can also be connected with Turcot syndrome. Additionally, the role of this gene in carcinogenic processes in other organs, especially in breast cancer, is also investigated (BRYŚ et al. 2004).

A study conducted by BAN and YANG (1998) has shown that MutL gene of *E. coli* is absolutely Mg-dependent and, in absence of magnesium ions, hydrolysis of the MutL-ATP complex can be observed. High homology between MHL and MutL suggests that magnesium is also indispensable for the activity of human MHL genes. Moreover, double-stranded DNA break repair induced by ionizing radiation or formed during meiosis has also been found to be Mg-dependent (HARTWIG 2001).


### Magnesium and enzymes

An activator of over 300 different enzymes, magnesium participates in many metabolic processes, including transformation of proteins, lipids, carbohydrates and nucleic acids as well as electrolyte transport across cell membranes (WOLF, CITTADINI 2003).


Magnesium can originally bind to a substrate (by chelation), producing a complex that is the correct substrate for enzyme or directly attach to enzyme, creating active structure able to affect a substrate. However, these mechanisms are combined with each other because ATPase affects the correct substrate (ATP-Mg) only if it is activated by another  $Mg^{2+}$  ion.

The general mechanisms of  $Mg^{2+}$  action as a cofactor can be described as follows (WOLF, CITTADINI 2003):


A. Magnesium is engaged in stabilization of an intermediate product:



B. Magnesium stabilizes a product leaving group:



C. Magnesium binds two reactive substrates simultaneously and facilitates reaction through the proximity effect:



The majority of enzymes can be also activated by metal ions other than  $\text{Mg}^{2+}$ , although such a replacement leads to reduced efficiency of enzymatic reaction.

### Magnesium in metabolic cycles

In higher organisms, metabolic processes such as glycolysis, Krebs cycle,  $\beta$ -oxidation, active transport of ions or electrochemical coupling are regulated by Mg-dependent enzymes. The main domain of magnesium action is the activation of enzymes responsible for formation, storing and using of high-energy compounds. All reactions involving ATP require the presence of magnesium ions (TOUYZ 2004). An  $\text{Mg}^{2+}$  ion, coupled with oxygen atoms of phosphorus groups located at  $\alpha$  and  $\beta$  positions, protects APT molecules from enzymatic hydrolysis, while the dislocation of  $\text{Mg}^{2+}$  in the direction of  $\alpha$  and  $\beta$  positions facilitates the hydrolysis of terminal phosphorus groups. Magnesium in the form of  $\beta$ ,  $\gamma$ -Mg-ATP complex binds to active centers of many enzymes. The complexes of Mg-ATP are essential for catalytic activity of, e.g., phosphotransferases (kinases), nucleotidylotransferases and ATPases (COWAN 1995).

Magnesium ions activating adenylate cyclase control cyclic adenosine monophosphate (cAMP) synthesis. Adenylate cyclase activation is crucial for the control of anaphylactic reactions because high intracellular cAMP and cGMP concentrations slow down or stop degranulation of mast cells. Consequently, the accessibility of magnesium to the enzyme can modulate cyclic nucleotide metabolisms in cells. Since  $\text{Mg}^{2+}$  deficit stimulates histamine release from mast cells by inhibition of cAMP production, it is believed that magnesium reduces the hypersensitivity reactions (BLACH et al. 2007).

Perfect confirmation of the key role of cellular magnesium is glycolysis, especially in human erythrocytes, as many enzymes involved in this process are Mg-dependent. Moreover, removing extracellular  $\text{Mg}^{2+}$  or chelating intracellular  $\text{Mg}^{2+}$  markedly inhibits glycolysis and limits glucose transport by erythrocytes (LAUGHLIN, THOMPSON 1996).

Numerous literature data suggest some correlation between glucose transport and changes in intro- or extracellular  $\text{Mg}^{2+}$  level resulting from hormonal stimulation of  $\beta$ -pancreatic islets (HENQUIN et al. 1983, FAGAN, ROMANI 2000), hepatocytes (FAGAN, SCARPA 2002, GAUSSIN et al. 1997) or cardiomyocytes (ROMANI et al. 1993, ROMANI, SCARPA 1990, ROMANI et al. 2000).

An increase in catecholamine or glucagon leads to secretion of glucose and  $Mg^{2+}$  from liver cells into the extracellular compartment (GAUSSIN et al. 1997). The presence of glucose transport inhibitors (ROMANI, SCARPA 1990) and the absence of extracellular  $Na^+$ , which hampers magnesium extrusion, also impair glucose output by liver cells. TORRES et al. (2005) have reported that hepatocytes from starved rats (after overnight fasting) accumulated approximately fourfold more  $Mg^{2+}$  than liver cells from fed animals. This clearly indicates that diminution of intrahepatic cellular glycogen or glucose level causes decreased ability of catecholamine or glucagon to mobilize  $Mg^{2+}$  from the hepatocyte.

In cardiac myocytes, parallel accumulation of glucose and  $Mg^{2+}$  is induced by insulin (HARTWIG 2001, ROMANI et al. 1993, 2000). Insulin acts as an endogenous regulating factor of  $Mg^{2+}$  homeostasis. Flux concentrations of blood or tissue magnesium are dependent on the amount of insulin released from pancreatic islets and insulin immunity of tissues (ROMANI et al. 2000). Also in this case, the absence of extracellular glucose or the presence of glucose transport inhibitors hamper  $Mg^{2+}$  transportation, lower extracellular  $Mg^{2+}$  content and break the transport of glucose to myocytes (ROMANI et al. 1993). Some correlation between glucose and  $Mg^{2+}$  transport/utilization in rats rendered diabetic by streptozotocin injection has been confirmed by FAGAN et al. (2004). Rats experienced a 10% and 20% decrease in the liver magnesium level after 4 and 8 weeks, respectively, after the onset of the disease. CEFARATTI et al. (2004) have confirmed diminished accumulation of  $Mg^{2+}$  in liver blisters isolated from rats with experimental diabetes.

Since  $Mg^{2+}$  accumulation directly or indirectly influences protein kinase C activation, it is possible that in diabetic patients the enzymatic action is disturbed. TANG et al. (1993) have also observed selective alterations in the expression PKC and marked differences in the distribution of the various isoforms between membrane and cytosol fractions of hepatocytes with streptozotocin-induced animals.

Similar modifications in the distribution of PKC as well as the reduction of cell magnesium content have been observed in tissues of ethanol-fed rats (YOUNG et al. 2003). The total magnesium concentration in animal hepatocytes of the examined group was  $26.8 \pm 2.4$  nM  $mg^{-1}$  proteins versus  $36.0 \pm 1.4$  nM  $mg^{-1}$  proteins for the control group. In comparison to the control conditions, the  $Mg^{2+}$  level in hepatocytes from EtOH-treated samples did not increase following stimulation of protein kinase C by vasopressin or analogs of diacylglycerol (DAG). Moreover, the stimulation of  $\alpha$ - or  $\beta$ -adrenoceptors in alcohol supplemented animals, did not elicit  $Mg^{2+}$  extrusion from liver cells to the extracellular space.

KIMURA et al. (1996) have shown that in Mg-deficient rats concentration of blood glucose and plasma insulin both in overnight fasted and non-fasted individuals as well as in response to oral sucrose loading are impaired. After 8 weeks of low- $Mg^{2+}$  diet, translocation of insulin-stimulated glucose trans-

porter 4 (GLUT4) to the adipocyte plasma membrane was significantly reduced. In addition, phosphorylation of insulin receptor was lower in Mg-deficient animals. On the other hand, wortmannin (WT) or another PI 3-kinase inhibitor blocked the insulin-stimulated activity of  $\text{Na}^+/\text{Mg}^{2+}$  exchange (FERREIRA et al. 2004).

These data may suggest that  $\text{Mg}^{2+}$  absence induces alterations in glucose metabolism by reducing intestinal glucose absorption or glucose assimilation in liver and/or other tissues (KIMURA et al. 1996).

### **Magnesium and cellular respiration**

Magnesium maintains a mitochondrial respiratory coupling chain, in which phosphorylation and oxidation obtain high efficiency. Magnesium ions might be transported to the mitochondrial matrix across Mrs2p channel of the inner mitochondrial membrane, whose activity depends on both electric potential and  $\text{Mg}^{2+}$  concentration, but the electrophysiological profile of Mrs2p remains to be developed (KOLISEK 2003).

The flux  $\text{Mg}^{2+}$  level in the mitochondrial matrix modulates  $\alpha$ -ketoglutarate dehydrogenase (CHAKRABORTI et al. 2002), pyruvate dehydrogenase and glutamate dehydrogenase (PANOV, SCARPA 1996) activity. Alterations of the matrix  $\text{Mg}^{2+}$  concentration (coupled with alternatively to changes of  $\text{Ca}^{2+}$ ) are reflected in the mitochondrial respiration rate.

Moreover, LIN et al. (1993) demonstrated that  $\text{Mg}^{2+}$  is an integral component of subunit IV of cytochrome c oxidase complex, the last enzyme of the respiratory chain catalyzing molecular oxygen reduction. The volume of an organelle is regulated by the matrix magnesium through direct control of the  $\text{K}^+/\text{H}^+$  antiporter, inhibition of mitochondrial inner membrane anion channel (IMAC) as well as through indirect modulation of the channel's permeability.

IMAC channels display selectivity among monovalent ( $\text{Cl}^-$ ,  $\text{HCO}_3^-$ ) as well as polyvalent (e.g. citrate) ions. These channels are probably also involved in the synchronization of oscillation in a mitochondrial membrane potential of isolated cardiomyocytes. Evidence has been provided that IMAC regulates the flow of anionic peroxidase from mitochondria during the ischaemic preconditioning (IPC) (SKALSKA et al. 2006). Although the IMAC control mechanism has not been completely elucidated, BEAVIS and POWERS (2004) suggested that the matrix  $\text{Mg}^{2+}$  as well as the protons impair channel activity.

The IMAC activation precedes the opening of the mitochondrial permeability transition pore (PTP), thereby promoting cell death. The PTP opening is a direct cause of the death of neurons in a damaged brain or in cardiac myocytes during ischemia and reperfusion. PTP also plays a role in muscular dystrophy (DMD), caused by deficiency of collagen VI, as well as in hepatocytosis, induced by cancerogenic factors (SKALSKA et al. 2006). The increase of the mitochondrial calcium pool facilitates the PTP opening whereas

a larger matrix  $Mg^{2+}$  concentration blocks this channel. Moreover, ZORATTI and SZABO (1995) showed that megachannels are inhibited by divalent cations, such as  $Mg^{2+}$  or  $Mn^{2+}$ , nucleotides: ADP and ATP as well as polyamines. DOLDER et al. (2003) established that magnesium plays an indirect role in modulating the PTP opening. They proved that creatine kinase can regulate the PTP size by tightly associating to the mitochondrial membrane and remaining in an active state. Impaired concentration of the extramitochondrial  $Mg^{2+}$  causes reduction of creatine kinase activity and increased pore permeability (DOLDER et al.).

Magnesium ions are also essential to glutathione synthesis, which can be confirmed by the fact that GSH level in the red blood cells of rats decreased after 2-3 weeks of a  $Mg^{2+}$ -deficient diet (WĘGLICKI et al. 1996). Glutathione depletion enforces reactive oxygen species accumulation, resulting in mitochondrial dysfunction, which is decisive in apoptotic cascade. The changes in the mitochondrial membrane's potential lead to the opening of megachannels in mitochondrial membranes, to alterations of membrane permeability, to translocation of cytochrome c and apoptosis inducing factor (AIF) from the mitochondria to the cytosol, which is the starting point for programmed cell death.

The above facts confirm the key role of  $Mg^{2+}$  in the regulation of mitochondrial function, including the control of their volume, composition of ions and ATP production.

### **Magnesium and calcium ion transport**

Magnesium ions are important for maintaining cell homeostasis because they are essential to the stabilization of cell membranes, to the activation of sodium-potassium pump (Na-K-ATP-ase) or calcium pump (Ca-ATP-ase), and to the regulation of composition of intra- and extracellular liquid (HARTWIG 2001, COWAN 1995).

As calcium antagonist, magnesium increases the neuromuscular excitability and has an antispastic and anticonvulsive effect, impairing the contractibility of muscles.

As early as in 1988, WHITE and HARTZELL showed that free intracellular magnesium can regulate the functioning of calcium channels. BARA and GUIET-BARA (2001) have confirmed that extracellular magnesium salts ( $MgCl_2$  or, to a smaller extent,  $MgSO_4$ ) reduce the influx of calcium through high-voltage channel  $Ca^{2+}$  type L in vascular smooth muscle cells (VSMCs) and vascular endothelial cells (VECs) of human placenta (BARA, GUIET-BARA 2001), and consequently modulate the tonus of placental vessels.  $Mg^{2+}$  and GTP binding sites are assumed to reside in the intracellular C-terminal side of the  $\alpha 1$  subunit of the channel. In basal conditions (i.e. the dephosphorylated channel and  $Mg^{2+}$  and GTP abundant on the intracellular side)  $Mg^{2+}$  and GTP binding to C-terminal inhibit the current conduction. A decrease in  $Mg^{2+}$  without intracellular GTP produces a current conducting state but

addition of GTP blocks the channel. Phosphorylation results in both  $Mg^{2+}$  and GTP blocks by unbinding these blocking substances through conformational change of the channel protein.

Serrano has described a similar blocking effect of extracellular magnesium on  $\alpha 1G$  T-type calcium channels, which play an important role in the mechanisms underlying thalamocortical oscillation (SERRANO et al. 2000). This is particularly essential because T channels are not blocked by classic calcium antagonists (except for mibefradil which is not used in clinical practice on account of undesirable action).

Whether or not  $Mg^{2+}$  ions modulate the action of store-operated calcium release-activated  $Ca^{2+}$  channels (CRAC), involved in regulation of inflammatory mediators production in allergic reactions as well as in differentiation and activation of T lymphocytes, is still not completely elucidated.

While the results of some experimental research have shown that intracellular magnesium modulates activity and selectivity of CRAC, others suggest that the channels regulated by intracellular  $Mg^{2+}$  are not CRAC channels but rather Mg-inhibited cation (MIC) channels that open as  $Mg^{2+}$  is washed out of the cytosol. MIC have been defined as another class of channels because they display different functional parameters from those displayed by CRAC in terms of inhibition (e.g. MIC are not blocked by SKF 96365 – the inhibitor of CRAC channels) or selectivity (unlike CRAC, MIC channels are permeable to  $Cs^+$  ions;  $PCs/PNa = 0.13$  vs.  $1.2$  for MIC) (PRAKRITYA, LEWIS 2000).

Studies carried out on rats with arterial hypertension have confirmed that extracellular  $Mg^{2+}$  imitates nifedipine in the process of reducing  $Ca^{2+}$  entry to vascular smooth muscle cells through store-operated channels (SOCs), resulting in the widening of circular vessels and a decrease in peripheral resistance as well as blood pressure (ZHANG et al. 2002).

## **Magnesium and potassium ion transport**

Potassium channels play a crucial role in the regulation of membrane potential in smooth muscle cells and vascular tone.

As the equilibrium potential for potassium ions in vascular smooth muscle cells is more negative ( $-84$  mV) than the cell's resting potential ( $-60$  to  $-70$  mV), the opening of potassium channels induces the K ion outflow from the cell. The loss of cations caused by an increase in the absolute value of membrane potential leads to the closing of L-type voltage-gated calcium channels (VGCC-L), to a decrease in intracellular calcium concentration as well as to relaxation of vessels. Blocking of potassium channels, however, lowers membrane potential, stimulates calcium ion inflow via voltage-gated ion channels (VDCC) and produces vessel contraction (BARANOWSKA et al. 2007).

TAMMARO et al. (2005) have provided evidence that intracellular  $Mg^{2+}$  ions affect voltage-dependent K channels ( $K_v$ ), which regulate potassium ion dis-

tribution and cooperate with  $K_{Ca}$  channels in control of arterial vessels con-  
volution in vascular smooth muscle cells. It was observed that an increase  
in the intracellular  $Mg^{2+}$  level slows down the  $K_V$  channel activation, caus-  
es inward rectification at positive membrane potentials and shifts voltage-  
dependent inactivation. The above results demonstrate that intracellular  
 $Mg^{2+}$  can act as a potent modulator of  $K_V$  channel in vascular smooth mus-  
cle cells, representing a novel mechanism for the regulation of  $K_V$  channel  
activity in the vasculature.

Cell magnesium also regulate the action of  $Ca^{+}$ -dependent  $K^{+}$  channels  
( $BK_{Ca}$ ), essential for modulating muscle contraction and neuronal activities  
such as synaptic transmission or hearing [http://www.nature.com/nature/jour-  
nal/v418/n6900/full/nature00941.html](http://www.nature.com/nature/journal/v418/n6900/full/nature00941.html) - B1 (SHI et al. 2002). Physiological ac-  
tivation of  $BK_{Ca}$  channels counteracts depolarization of cell membranes, con-  
traction of blood vessels and increasing pressure (BARANOWSKA et al. 2007).  
Because of the importance of BK channels in neurotransmitter release and  
vascular tone,  $Mg^{2+}$  modulation of BK channels may play a substantial role  
in these pathophysiological processes.  $Mg^{2+}$  modulates their permeability by  
blocking the opening of a BK channel or by stimulation of channels inde-  
pendently from  $Ca^{2+}$  and voltage changes following binding to an open chan-  
nel in different than Ca specific site or in no site (SHI et al. 2002). The  
structural separation between the binding site and the activation gate indi-  
cates that  $Mg^{2+}$  binding activates the channel by an allosteric mechanism;  
i.e.,  $Mg^{2+}$  binding may cause a conformational change at the binding site  
that propagates to the activation gate for a channel opening (HUANGHE 2008).

Intracellular  $Mg^{2+}$  affects bioelectrical activity of the heart via regula-  
tion of inward rectifying potassium channels ( $K_{IR}$ ), which are responsible for  
blocking outflow of K ions from cell and repolarization (BARANOWSKA et al.  
2007).

Physiological concentrations of intracellular Mg-ADP complex regulate  
the sensitivity of ATP-sensitive potassium channels ( $K_{ATP}$ ) to sulphonylurea  
derivatives. Sulphonylurea derivatives, used to treat type 2 diabetes, stimu-  
late insulin secretion by blocking  $K_{ATP}$  channels in pancreatic  $\beta$ -cells. An  
intracellular Mg-ADP complex modulates sulphonylurea block, enhancing the  
inhibition of Kir6.2/SUR1 ( $\beta$ -cell type) and decreasing that of Kir6.2/SUR2A  
(cardiac-type) channels. This is important because the opening of  $K_{ATP}$  chan-  
nels is regarded as an endogenous cardioprotective mechanism so the block-  
ing effect of sulphonylurea derivatives in the cardiovascular system may  
have deleterious effects (REIMANN et al. 2003).

The influence of  $Mg^{2+}$  on  $K^{+}$  channels is not limited to the cell mem-  
brane. BEDNARCZYK et al. (2005) have shown that matrix  $Mg^{2+}$  ions affect  
mitochondrial ATP-dependent potassium channel ( $K_{ATP}$ ) in the heart, which  
plays a key role in protecting from ischemia/reperfusion. The ATP/ $Mg^{2+}$  com-  
plex inhibits  $K_{ATP}$  activity and free magnesium ions regulate both the chan-  
nel conductance and open probability. Another study has suggested that mi-

to  $K_{ATP}$  channels make functional connection with mitochondrial pyruvate dehydrogenase forming a larger, multiprotein complex. A hypothesis has been formulated that enhanced activity of  $mitoK_{ATP}$  channel protects the heart muscle during myocardial ischemia as well as neurons, brain cells and skeletal muscle cells (SKALSKA et al. 2006).

## REFERENCES

- ADAMALA K., PIKUŁA S. 2004. *Hipotetyczna rola autokatalitycznych właściwości kwasów nukleinowych w procesie biogenezy. [A hypothetical role of autocatalytic properties of nucleic acids in biogenesis]*. Kosmos, 53 (2): 123-131 (in Polish).
- ANASTASSOPOULOU J., THEOPHANIDES T. 2002. *Magnesium - /DNA interactions and the possible relation of magnesium to carcinogenesis. Irradiation and free radicals*. Crit. Rev. Oncol./Hematol., 42 (1): 79-91.
- BAN C., YANG W. 1998. *Crystal structure and ATPase activity of MutL: implications for DNA repair and mutagenesis*. Cell, 95: 541-522.
- BARA M., GUIET-BARA A. 2001. *Magnesium regulation of  $Ca^{2+}$  channels in smooth muscle and endothelial cells of human allantochorial placental vessels*. Magnes. Res., 14: 11-18.
- BARANOWSKA M., KOZŁOWSKA H., KORBUĆ A. et al. 2007. *Kanały potasowe w naczyniach krwionośnych – ich znaczenie w fizjologii i patologii. [Potassium channels in blood vessels: Their role in health and disease]*. Post. Hig. Med. Dośw., 61: 596-605 (in Polish).
- BEAVIS D., POWERS M. 2004. *Temperature dependence of the mitochondrial inner membrane anion channel*. J. Biol. Chem., 279: 4045-4050.
- BEDNARCZYK P., DOŁOWY K., SZEWCZYK A. 2005. *Matrix  $Mg^{2+}$  regulates mitochondrial ATP-dependent potassium channel from heart*. FEBS Lett., 579: 1625-1632.
- BŁACH J., NOWACKI W., MAZUR A. 2007. *Wpływ magnezu na reakcje alergiczne skóry. [Magnesium in skin allergy]*. Post. Hig. Med. Dośw., 61: 548-557. (in Polish)
- BRYŚ M., KRAJEWSKA W.M., ZYCH A. et al. 2004. *Mutacje genu hMLH1 a sporadyczny rak piersi kobiet. [Mutations of hMLH1 gene and sporadic breast cancer]* Prz. Menopauz., 6: 47-50 (in Polish).
- CEFARATTI CH., MCKINNIS A., ROMANI A. 2004. *Altered  $Mg^{2+}$  transport across liver plasma membrane from streptozotocin-treated rats*. Moll. Cell. Biochem., 262: 145-154.
- CERIELLO A., MOTZ A. 2004. *Is oxidative stress the pathogenic mechanism underlying insulin resistance, diabetes and cardiovascular disease? The common soil hypothesis revisited*. Arterioscler. Thromb. Vasc. Biol., 24: 816-823.
- CHAKRABORTI S., CHAKRABORTI T., MANDAL M. et al. 2002. *Protective role of magnesium in cardiovascular diseases: A review*. Mol. Cell. Biochem., 238: 163-179.
- COOK J.A. 2000. *The genetics and management of inherited gynaecological cancer (including breast)*. Curr. Obstet. Gynaecol., 10: 133-138.
- COWAN J.A. 1995. *Introduction to the biological chemistry of magnesium ion. The biological chemistry of magnesium*. VCH. New York, 1-23.
- DOLDER M., WALZEL B., SPEER O. et al. 2003. *Inhibition of the mitochondrial permeability transition by creatine kinase substrates. Requirement for microcompartmentation*. J. Biol. Chem., 278: 17760-17766.
- FAGAN T.E., CEFARATTI CH., ROMANI A. 2004. *Streptozotocin-induced diabetes impairs  $Mg^{2+}$  homeostasis and uptake in rat liver cells*. Am. J. Physiol. Endocrinol. Metab., 286: 184-193.
- FAGAN T.E., ROMANI A. 2000. *Activation of  $Na^{+}$ - and  $Ca^{2+}$ -dependent  $Mg^{2+}$  extrusion by  $\alpha_1$ - and  $\beta$ -adrenergic agonists in rat liver cells*. Am. J. Physiol. Gastrointest. Liver Physiol., 279: 943-950.

- FAGAN T.E., SCARPA A. 2002. *Hormone-stimulated Mg<sup>2+</sup> accumulation into rat hepatocytes: a pathway for rapid Mg<sup>2+</sup> and Ca<sup>2+</sup> redistribution.* Arch. Biochem. Biophys., 401: 277-282.
- FERREIRA A., RIVERA A., ROMERO J.R. 2004. *Na<sup>+</sup>/Mg<sup>2+</sup> exchange is functionally coupled to the insulin receptor.* J. Cell. Physiol., 199 (3): 434-440.
- GAUSSIN V., GAILLY P., GILLIS J.M. et al. 1997. *Fructose-induced increase in intracellular free Mg<sup>2+</sup> ion concentration in rat hepatocytes: relation with the enzymes of glycogen metabolism.* Biochem. J., 326: 823-827.
- HAMPEL A., COWAN J.A. 1997. *A unique mechanism for RNA catalysis: the role of metal cofactors in hairpin ribozyme cleavage.* Chem. Biol., 4: 513-517.
- HARTWIG A. 2001. *Role of magnesium in genomic stability.* Mutat. Res., 475: 113-121.
- HENQUIN J.C., TAMAGAWA T., NENQUIN M. et al. 1983. *Glucose modulates Mg<sup>2+</sup> fluxes in pancreatic islet cells.* Nature, 301: 73-74.
- HUANGHE Y., LEI H., JINGYI S. et al. 2008. *Tuning magnesium sensitivity of BK channels by mutations.* Biophys. J., 91: 2892-2900.
- JOŚKO J., KNEFEL K. 2003. *The role of vascular endothelial growth factor in cerebral oedema formation.* Pol. Neuropathol., 43: 161-166.
- KIMURA Y., MURASE M., NAGATA Y. 1996. *Change in glucose homeostasis in rats by long-term magnesium-deficient diet.* J. Nutr. Sci. Vitaminol., 42: 407-422.
- KOLISEK M., ZSURKA G., SAMAJ J. et al. 2003. *Mrs2p is an essential component of the major electrophoretic Mg<sup>2+</sup> influx system in mitochondria.* EMBO J., 22: 1235-1244.
- LAUGHLIN M.R., THOMPSON D. 1996. *The regulatory role for magnesium in glycolytic flux of the human erythrocyte.* J. Biol. Chem., 271: 28977-28983.
- LIN J., PAN L.P., CHAN S.I. 1993. *The subunit location of magnesium in cytochrome c oxidase.* J. Biol. Chem., 268: 22210-22214.
- PANOV A., SCARPA A. 1996. *Mg<sup>2+</sup> control of respiration in isolated rat liver mitochondria.* Biochemistry, 35 (39): 12849-12856.
- PRAKRIYA M., LEWIS R.S. 2000. *Separation and characterization of currents through store-operated CRAC channels and Mg-inhibited cation (MIC) channels.* J. Gen. Physiol., 119 (5): 487-507.
- REIMANN F., DABROWSKI M., JONES P. et al. 2003. *Analysis of the differential modulation of sulphonylurea block of  $\beta$ -cell and cardiac ATP-sensitive K<sup>+</sup> (K<sub>ATP</sub>) channels by Mg-nucleotides.* J. Physiol., 547: 159-168.
- ROMANI A., MARFELLA C., SCARPA A. 1993. *Cell magnesium transport and homeostasis: role of intracellular compartments.* Miner. Electrol. Metab., 19: 282-289.
- ROMANI A., MATTHEWS V., SCARPA A. 2000. *Parallel stimulation of glucose and Mg<sup>2+</sup> accumulation by insulin in rat hearts and cardiac ventricular myocytes.* Circ. Res., 86: 326-333.
- ROMANI A., SCARPA A. 1990. *Hormonal control of Mg<sup>2+</sup> transport in the heart.* Nature, 346: 841-844.
- ROSZKOWSKI K. 2002. *Mechanizmy naprawy oksydacyjnych uszkodzeń DNA. [Repair mechanisms of oxidative DNA damage]* Współcz. Onkol., 6 (6): 360-365 (in Polish).
- SANCAR A. 1994. *Mechanisms of DNA excision repair.* Science, 266: 1994-1996.
- SERRANO J.R., DASHTI S.R., PEREZ-REYES E. et al. 2000. *Mg<sup>2+</sup> block unmasks Ca<sup>2+</sup>/Ba<sup>2+</sup> selectivity of  $\alpha$ 1G T-type calcium channels.* Biophys. J., 79: 3052-3062.
- SHI J., KRISHNAMOORTHY G., YANG Y. et al. 2002. *Mechanism of magnesium activation of calcium-activated potassium channels.* Nature, 418: 876-880.
- SKALSKA J., DĘBSKA-VIELHABER G., GŁAB M. et al. 2006. *Mitochondrialne kanały jonowe. [Mitochondrial ion channels].* Post. Biochem, 52 (2): 137-144 (in Polish).

- TAMMARO P., SMITH A.L., CROELEY B.L. et al. 2005. *Modulation of the voltage-dependent K<sup>+</sup> current by intracellular Mg<sup>2+</sup> in rat aortic smooth muscle cells.* Cardiovasc. Res., 65: 387-396.
- TANG E.Y., PARKER P.J., BEATTIE J. et al. 1993. *Diabetes induces selective alterations in the expression of protein kinase C isoforms in hepatocytes.* FEBS Lett., 326: 117-123.
- TORRES L.M., YOUNGNER J., ROMANI A. 2005. *Role of glucose in modulating Mg<sup>2+</sup> homeostasis in liver cells from starved rats.* Am. J. Physiol. Gastrointest. Liver Physiol., 288: 195-206.
- TOUYZ R.M. 2004. *Magnesium in clinical medicine.* Front. Biosci., 9: 1278-1293.
- WEDEKIND J.E., REED G.H., RAYMENT I. 1995. *Octahedral coordination at the high-affinity metal site in enolase: crystallographic analysis of the MgII – enzyme complex from yeast at 1.9 Å resolution.* Biochemistry, 34 (13): 4325-4330.
- WĘGLICKI W.B., MAK I.T., KRAMER J.H. et al. 1996. *Role of free radicals and substance P in magnesium deficiency.* Cardiovasc. Res., 31: 677-687.
- WHITE R.E., HARTZELL H.C. 1988. *Effects of intracellular free magnesium on calcium current in isolated cardiac myocytes.* Science, 239: 778-780..
- WOLF F.I., CITTADINI A. 2003. *Chemistry and biochemistry of magnesium.* Mol. Asp. Med., 24: 3-9.
- WOLF F.I., TORSSELLO A., FANSANELLA S. et al. 2003. *Cell physiology of magnesium.* Mol. Asp. Med., 24: 11-26.
- YOUNG A., CEFARATTI CH., ROMANI A. 2003. *Chronic EtOH administration alters liver Mg<sup>2+</sup> homeostasis.* Am. J. Physiol. Gastrointest. Liver Physiol., 284 (1): 57-67.
- ZHANG J., WIER W.G., BLAUSTEIN M.P. 2002. *Mg<sup>2+</sup> blocks myogenic tone but not K<sup>+</sup>-induced constriction: role for SOCs in small arteries.* Am. J. Physiol. Heart Circ. Physiol., 283, 2692-2705.
- ZORATTI M., SZABO L. 1995. *The mitochondrial permeability transition.* Biochim. Biophys. Acta., 1241: 139-176.

---

**Reviewers of the Journal of Elementology Vol. 15(3), Y. 2010**

Wiesław Barabasz, Dariusz Bednarek, Wiesław Bednarek, Bolesław Bieniek,  
Janina Gajc-Wolska, Kazimierz Grabowski, Stanisław Ignatowicz,  
Maria Iskra, Adam Kaczor, Eugeniusz Kołota, Ireneusz Kowalski,  
Aleksandra Kwiatkowska, Zenia Michałojć, Andrzej Sapek,  
Małgorzata Schleger-Zawadzka, Lech Walasek, Czesław Wołoszyk



## Regulamin ogłaszania prac w „Journal of Elementology”

1. Journal of Elementology (kwartalnik) zamieszcza na swych łamach prace oryginalne, doświadczalne, kliniczne i przeglądowe z zakresu przemian biopierwiastków i dziedzin pokrewnych.
2. W JE mogą być zamieszczone artykuły sponsorowane, przygotowane zgodnie z wymaganiami stawianymi pracom naukowym.
3. W JE zamieszczamy materiały reklamowe.
4. Materiały do wydawnictwa należy przesłać w 2 egzemplarzach. Objętość pracy oryginalnej nie powinna przekraczać 10 stron znormalizowanego maszynopisu (18 000 znaków), a przeglądowej 15 stron (27 000 znaków).
5. Układ pracy w języku angielskim: TYTUŁ PRACY, imię i nazwisko autora (-ów), nazwa jednostki, z której pochodzi praca, streszczenie w języku angielskim i polskim – minimum 250 słów. Streszczenie powinno zawierać: wstęp (krótko), cel badań, metody badań, omówienie wyników, wnioski. Przed streszczeniem w języku angielskim: Abstract (tekst streszczenia), Key words (maks. 10 słów). Przed streszczeniem w języku polskim: TYTUŁ PRACY, Abstrakt, (tekst streszczenia), Słowa kluczowe: (maks. 10 słów). WSTĘP, MATERIAŁ I METODY, WYNIKI I ICH OMÓWIENIE, WNIOSKI, PIŚMIENNICTWO. U dołu pierwszej strony należy podać tytuł naukowy lub zawodowy, imię i nazwisko autora oraz dokładny adres przeznaczony do korespondencji w języku angielskim.
6. Praca powinna być przygotowana wg zasad pisowni polskiej. Jednostki miar należy podawać wg układu SI, np.: mmol(+) kg<sup>-1</sup>; kg ha<sup>-1</sup>; mol dm<sup>-3</sup>; g kg<sup>-1</sup>; mg kg<sup>-1</sup> (obowiązują formy pierwiastkowe).
7. W przypadku stosowania skrótu po raz pierwszy, należy podać go w nawiasie po pełnej nazwie.
8. Tabele i rysunki należy załączyć w oddzielnych plikach. U góry, po prawej stronie tabeli należy napisać Tabela i numer cyfrą arabską, również w języku angielskim, następnie tytuł tabeli w języku polskim i angielskim wyrównany do środka akapitu. Ewentualne objaśnienia pod tabelą oraz opisy tabel powinny być podane w języku polskim i angielskim. Wartości liczbowe powinny być podane jako zapis złożony z 5 znaków pisarskich (np. 346,5; 46,53; 6,534; 0,653).
9. U dołu rysunku, po lewej stronie, należy napisać Rys. i numer cyfrą arabską oraz umieścić podpisy i ewentualne objaśnienia w języku polskim i angielskim.
10. Piśmiennictwo należy uszeregować alfabetycznie, bez numerowania, w układzie: NAZWISKO INICJAŁ IMIENIA (KAPITALIKI), rok wydania. Tytuł pracy (kursywa). Obowiązujący skrót czasopisma, tom (zeszyt): strony od-do, np. KOWALSKA A., KOWALSKI J. 2002. *Zwartość magnezu w ziemiakach*. Przem. Spoż., 7(3): 23-27. Tytuły publikacji wyłącznie w języku angielskim z podaniem oryginalnego języka publikacji, np. (in Polish).
11. W JE można także cytować prace zamieszczone w czasopismach elektronicznych wg schematu: NAZWISKO INICJAŁ IMIENIA (KAPITALIKI), rok wydania. Tytuł pracy (kursywa). Obowiązujący skrót czasopisma internetowego oraz pełny adres strony internetowej. np. ANTONKIEWICZ J., JASIEWICZ C. 2002. *The use of plants accumulating heavy metals for detoxication of chemically polluted soils*. Electr. J. Pol. Agric. Univ., 5(1): 1-13. hyperlink "<http://www.ejpau.media.pl/series/volume5/issue1/environment/art-01.html>
12. W pracach naukowych nie cytujemy podręczników, materiałów konferencyjnych, prac nierecenzowanych, wydawnictw popularnonaukowych.
13. Cytując piśmiennictwo w tekście, podajemy w nawiasie nazwisko autora i rok wydania pracy (KOWALSKI 1992). W przypadku cytowania dwóch autorów, piszemy ich nazwiska rozdzielone przecinkiem i rok (KOWALSKI, KOWALSKA 1993). Jeżeli występuje większa liczba nazwisk, podajemy pierwszego autora z dodatkiem i in., np.: (KOWALSKI i in. 1994). Cytując jednocześnie kilka pozycji, należy je uszeregować od najstarszej do najnowszej, np.: (NOWAK 1978, NOWAK i in. 1990, NOWAK, KOWALSKA 2001).

14. Do artykułu należy dołączyć pismo przewodnie Kierownika Zakładu z jego zgodą na druk oraz oświadczenie Autora (-ów), że praca nie została i nie zostanie opublikowana w innym czasopiśmie bez zgody Redakcji JE.
15. Dwie kopie wydruku komputerowego pracy (Times New Roman 12 pkt przy odstępie akapitu 1,5 - bez dyskietki) należy przesłać na adres Sekretarzy Redakcji:

**dr hab. Jadwiga Wierzbowska, prof. UWM**  
**Uniwersytet Warmińsko-Mazurski w Olsztynie**  
**Katedra Chemii Rolnej i Ochrony Środowiska**  
**ul. Oczapowskiego 8, 10-719 Olsztyn-Kortowo**  
**jadwiga.wierzbowska@uwm.edu.pl**

**dr hab. Katarzyna Glińska-Lewczuk**  
**University of Warmia and Mazury in Olsztyn**  
**Pl. Łódzki 2, 10-759 Olsztyn, Poland**  
**kaga@uwm.edu.pl**

16. Redakcja zastrzega sobie prawo dokonywania poprawek i skrótów. Wszelkie zasadnicze zmiany tekstu będą uzgadniane z Autorami.
17. Po recenzji Autor zobowiązany jest przesłać w 2 egzemplarzach poprawiony artykuł wraz z nośnikiem elektronicznym (dyskietka, CD lub e-mail), przygotowany w dowolnym edytorze tekstu, pracującym w środowisku Windows.

**Redakcja Journal of Elementology uprzejmie informuje:**

Koszt wydrukowania maszynopisu (wraz z rysunkami, fotografiami i tabelami) o objętości nieprzekraczającej 6 stron formatu A4, sporządzonego wg następujących zasad:

- czcionka: Times New Roman, 12 pkt, odstęp 1,5;
  - 34 wiersze na 1 stronie;
  - ok. 2400 znaków (bez spacji) na 1 stronie;
  - rysunki i fotografie czarno-białe;
- wynosi 250 PLN + VAT.

Koszt druku każdej dodatkowej strony (wraz z rysunkami, fotografiami i tabelami) wynosi 35 PLN + VAT.

Koszt druku 1 rysunku lub fotografii w kolorze wynosi 150 PLN + VAT.

**Uwaga:**

Z opłaty za druk pracy zwolnieni zostaną lekarze niezatrudnieni w instytutach naukowych, wyższych uczelniach i innych placówkach badawczych.

**Warunki prenumeraty czasopisma:**

- członkowie indywidualni PTMag 40.00 PLN + 0% VAT rocznie,
- osoby fizyczne 50.00 PLN + 0% VAT rocznie,
- biblioteki i instytucje 150 PLN + 0% VAT rocznie za 1 komplet (4 egzemplarze)  
+ 10.00 PLN + VAT za przesyłkę

Wpłaty prosimy kierować na konto UWM w Olsztynie:  
PKO S.A. I O/Olsztyn, 32124015901111000014525618  
koniecznie z dopiskiem "841-2202-1121"

## Guidelines for Authors „Journal of Elementology”

1. Journal of Elementology (a quarterly) publishes original scientific or clinical research as well as reviews concerning bioelements and related issues.
2. Journal of Elementology can publish sponsored articles, compliant with the criteria binding scientific papers.
3. Journal of Elementology publishes advertisements.
4. Each article should be submitted in duplicate. An original paper should not exceed 10 standard pages (18 000 signs). A review paper should not exceed 15 pages (27 000 signs).
5. The paper should be laid out as follows: **TITLE OF THE ARTICLE, name and surname of the author(s), the name of the scientific entity, from which the paper originates, INTRODUCTION, MATERIAL AND METHODS, RESULTS AND DISCUSSION, CONCLUSIONS, REFERENCES**, abstract in the English and Polish languages, min. 250 words. Summary should contain: introduction (shortly), aim, results and conclusions. Prior to the abstract in the English language the following should be given: **name and surname of the author(s), TITLE**, Key words (max 10 words), Abstract, **TITLE**, Key words and Abstract in Polish. At the bottom of page one the following should be given: scientific or professional title of the author, name and surname of the author, detailed address for correspondence in the English and Polish languages.
6. The paper should be prepared according to the linguistic norms of the Polish and English language. Units of measurements should be given in the SI units, for example  $\text{mmol}(+)\cdot\text{kg}^{-1}$ ;  $\text{kg ha}^{-1}$ ;  $\text{mol dm}^{-3}$ ;  $\text{g kg}^{-1}$ ;  $\text{mg kg}^{-1}$  (elemental forms should be used).
7. In the event of using an abbreviation, it should first be given in brackets after the full name.
8. Tables and figures should be attached as separate files. At the top, to the right of a table the following should be written: Table and table number in Arabic figures (in English and Polish), in the next lines the title of the table in English and Polish adjusted to the centre of the paragraph. Any possible explanation of the designations placed under the table as well as a description of the table should be given in English and Polish. Numerical values should consist of five signs (e.g. 346.5, 46.53, 6.534, 0.653).
9. Under a figure, on the left-hand side, the following should be written: Fig. and number in Arabic figures, description and possible explanation in Polish and English.
10. References should be ordered alphabetically but not numbered. They should be formatted as follows: Surname First Name Initial (capital letter) year of publication, Title of the paper (italics). The official abbreviated title of the journal, volume (issue): pages from – to. e.g. KOWALSKA A., KOWALSKI J. 2002. *Zawartość magnezu w ziemiakach*. Przem. Spoż., 7(3): 23-27.
11. It is allowed to cite papers published in electronic journals formatted as follows: Surname First Name Initial (capital letters) year of publication. Title of the paper (italics). The official abbreviated title of the electronic journal and full address of the website. e.g. ANTONKIEWICZ J., JASIEWICZ C. 2002. *The use of plants accumulating heavy metals for detoxication of chemically polluted soils*. Electr. J. Pol. Agric. Univ., 5(1): 1-13. hyperlink „<http://www.ejpau.pl/series/volume5/issue1/environment/art-01.html>” <http://www.ejpau.pl/series/volume5/issue1/environment/art-01.html>
12. In scientific papers, we do not cite textbooks, conference proceedings, non-reviewed papers and popular science publications.
13. In the text of the paper a reference should be quoted as follows: the author's name and year of publication in brackets, e.g. (KOWALSKI 1992). When citing two authors, their surnames should be separated with a comma, e.g. (KOWALSKI, KOWALSKA 1993). If there are more than two authors, the first author's name should be given followed

by et al., e.g. (KOWALSKI et al. 1994). When citing several papers, these should be ordered chronologically from the oldest to the most recent one, e.g. (NOWAK 1978, NOWAK et al. 1990, NOWAK, KOWALSKA 2001).

14. A paper submitted for publication should be accompanied by a cover letter from the head of the respective institute who agrees for the publication of the paper and a statement by the author(s) confirming that the paper has not been and will not be published elsewhere without consent of the Editors of the Journal of Elementology.
15. Two computer printed copies of the manuscript (Times New Roman 12 fonts, 1.5-spaced, without a diskette) should be submitted to the Editor's Secretary:

**dr hab. Jadwiga Wierzbowska, prof. UWM**  
**University of Warmia and Mazury in Olsztyn**  
**ul. Michała Oczapowskiego 8, 10-719 Olsztyn**  
**[jawierz@uwm.edu.pl](mailto:jawierz@uwm.edu.pl)**

**dr hab. Katarzyna Glińska-Lewczuk**  
**University of Warmia and Mazury in Olsztyn**  
**pl. Łódzki 2, 10-759 Olsztyn, Poland**  
**[kaga@uwm.edu.pl](mailto:kaga@uwm.edu.pl)**

16. The Editors reserve the right to correct and shorten the paper. Any major changes in the text will be discussed with the Author(s).
17. After the paper has been reviewed and accepted for publication, the Author is obliged to send the corrected version of the article together with the diskette. The electronic version can be prepared in any word editor which is compatible with Windows software.